

AD-A074 706

SCHOOL OF AEROSPACE MEDICINE BROOKS AFB TX
COMPARISON OF U.S.P. AND NON-U.S.P. GRADE IODINE-131: A PRELIMI--ETC(U)
AUG 79 D HUNTER, R E TATSCH

F/G 6/18

UNCLASSIFIED

SAM-TR-79-24


NL

1 OF 1
ADA
074706



END
DATE
FILMED

11 -79
DDC



AD A074706

LEVEL II

2
B.S.

Report SAM-TR-79-24

**COMPARISON OF U.S.P. AND NON-U.S.P. GRADE
IODINE-131: A PRELIMINARY REPORT**

**David Hunter, Captain, USAF, BSC
Robert E. Tatsch, B.S.**

August 1979

Interim Report for Period February 1979 - April 1979

Approved for public release; distribution unlimited.

DDC
RECEIVED
OCT 5 1979
D

DDC FILE COPY

**USAF SCHOOL OF AEROSPACE MEDICINE
Aerospace Medical Division (AFSC)
Brooks Air Force Base, Texas 78235**



79 09 24 347

NOTICES

This interim report was submitted by personnel of the Health Physics Branch, Radiation Sciences Division, USAF School of Aerospace Medicine, Aerospace Medical Division, AFSC, Brooks Air Force Base, Texas, under job order SUPTXHPS.

When U.S. Government drawings, specifications, or other data are used for any purpose other than a definitely related Government procurement operation, the Government thereby incurs no responsibility nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise, as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use, or sell any patented invention that may in any way be related thereto.

This report has been reviewed by the Information Office (OI) and is releasable to the National Technical Information Service (NTIS). At NTIS, it will be available to the general public, including foreign nations.

This technical report has been reviewed and is approved for publication.

David Hunter
DAVID HUNTER, Captain, USAF, BSC
Project Scientist

Robert E. Tatsch
ROBERT E. TATSCH, B.S.
Chief, Health Physics Branch

L. J. Enders
LAWRENCE J. ENDERS
Colonel, USAF, MC
Commander

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
Distribution/	
Availability Codes	
Avail and/or	special
A	

DDC
RECEIVED
OCT 5 1979
D

79 09 24 347

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER SAM-TR-79-24	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) COMPARISON OF U.S.P. AND NON-U.S.P. GRADE IODINE-131: A PRELIMINARY REPORT	5. TYPE OF REPORT & PERIOD COVERED Interim Report Feb 79 - Apr 79	6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) David Hunter Robert E. Tatsch	8. CONTRACT OR GRANT NUMBER(s)	
9. PERFORMING ORGANIZATION NAME AND ADDRESS USAF School of Aerospace Medicine (RZH) Aerospace Medical Division (AFSC) Brooks AFB, TX 78235	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 62202F SUPTXHPS	
11. CONTROLLING OFFICE NAME AND ADDRESS USAF School of Aerospace Medicine (RZH) Aerospace Medical Division (AFSC) Brooks AFB, TX 78235	12. REPORT DATE August 1979	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	13. NUMBER OF PAGES 2	15. SECURITY CLASS. (of this report) Unclassified
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Airborne radioactivity, airborne concentrations, volatilization of I-131		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) This report compares the airborne radioactivity from a U.S.P. grade and a non-U.S.P. grade Iodine-131 liquid sample. The results from this report indicate that the non-U.S.P. grade I-131 has a significantly higher airborne radioactivity level than the U.S.P. grade I-131.		

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

COMPARISON OF U.S.P. AND NON-U.S.P. GRADE IODINE-131: A PRELIMINARY REPORT

INTRODUCTION

A significant thyroid uptake in individuals who open and prepare oral solutions of Iodine-131 for therapeutic doses has been reported (1). The uptake and concentration of the I-131 in the individual's thyroid gland appears to be due to breathing volatile iodine which is released when the cap of the container is removed. Pollock and Myser (2) reported iodine releases in the hospital elevator and hallways as I-131 was being transported from the nuclear pharmacy to the patient's room. This preliminary report compares the airborne radioactivity from millicurie quantities of U.S.P. grade and non-U.S.P. grade I-131 in the liquid form.

MATERIALS AND METHODS

A carbon-impregnated cellulose filter was counted with an Ortec model 6200 multichannel analyzer and a 1 3/4" X 2" sodium iodide (Tl) crystal. The air sampler used for this study was a Dupont model 4000 with a calibrated air flow of 4 liters per minute. The air sampler efficiency and calculations of airborne radioactivity were performed in the same manner as by Browning et al. (1). The vial containing the I-131 was positioned under the holder of the air sampler. With the vial cap removed, the air sampler was allowed to run for 10 minutes. The distance from the iodine vials was the same for all measurements. Table 1 summarizes the liquid I-131 sample results.

TABLE 1. U.S.P. AND NON-U.S.P. GRADE I-131

<u>I-131</u>	<u>Total activity (mCi)</u>	<u>Volume (ml)</u>	<u>Concentration (mCi/ml)</u>	<u>Airborne radioactivity (μCi/ml)</u>
U.S.P.*	70.5	10	7.05	---
U.S.P.*	70.5	10	7.05	1.18×10^{-10}
U.S.P.*	70.5	10	7.05	1.13×10^{-9}
Non U.S.P.*	100.0	4	25.00	2.11×10^{-7}
Non U.S.P.*	38.0	2	19.00	7.03×10^{-9}

*E.R. Squibb & Sons

+Mallinckrodt, Inc.

DISCUSSION

As can be seen from Table 1, the airborne concentration with the non-U.S.P. iodine is significant. The airborne radioactivity from the 100 mCi non-U.S.P. grade iodine was approximately 29 times greater than that from the 38 mCi--which did not exceed the airborne limits set by the Nuclear Regulatory Commission at 9×10^{-9} $\mu\text{Ci/ml}$. Although limited by a maximum activity of 70.5 mCi, all of the U.S.P. grade iodine was well below these airborne radioactivity limits.

The data from this preliminary report indicate that the U.S.P. grade iodine has significantly less airborne radioactivity levels than the non-U.S.P. grade iodine. The non-U.S.P. grade iodine has a pH range between 2.0 and 4.0, while the U.S.P. grade iodine has a pH range between 7.5 and 9.0. Buffering the iodine solution at a pH between 7.5 and 9.0 maintains the preparation as sodium iodide, thereby preventing the volatilization of ^{131}I -hydriodic acid (3). Radiation hazards of personnel would be reduced by using U.S.P. grade iodine when therapeutic doses are required. The followup to this report will be conducted when the evaluation of higher activity levels of the U.S.P. grade I-131 is completed, both from the container and from breath samples of patients treated with therapeutic doses of 100 mCi or more of I-131.

REFERENCES

1. Browning, E. J., et al. Airborne concentration of I-131 in a nuclear medicine laboratory. Nucl Med 19:1078-1081 (1978).
2. Pollock, R. W., and R. D. Myser. Concentration of I-131 in the air during thyroid therapies. Health Phys 36(1):69-70 (1979).
3. Stafford, R. E. R. Squibb and Sons, San Antonio, Texas. Personal communication, 1979.